



## Horticultural and broadacre pesticides known to be toxic to honeybees in Australia – alphabetically sorted by product name

Product name	Active constituent(s) and concentration	Bee related label statement
4FARMERS ABAMECTIN 18 EC MITICIDE - INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
4FARMERS ALPHA-CYPERMETHRIN 100 EC INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
4FARMERS BIFENTHRIN 100 EC INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
4FARMERS CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
4FARMERS CYPERMETHRIN 260 EC INSECTICIDE	cypermethrin(260g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
4FARMERS DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
4FARMERS IMIDACLOPRID 200SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
4FARMERS OMETHOATE 290 INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AAKO RIDACIDE 400 EC INSECTICIDE	methidathion(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AAKOMECTINE MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ACRAMITE MITICIDE	bifenazate(480g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ACTARA INSECTICIDE	thiamethoxam (250g/kg)	This product is dangerous to bees and will kill any bees which are foraging in the crop to be treated or in hives which are oversprayed.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
ADONIS 3UL INSECTICIDE	fipronil(3g/L)	Dangerous to bees. DO NOT apply while bees are foraging. Ensure bees are not foraging in the crop to be treated. Remove beehives from areas due to be sprayed. Ensure alternative (untreated) source of nectar is available for bees.
ADONIS 8.5UL INSECTICIDE	fipronil(8.5g/L)	Dangerous to bees. DO NOT apply while bees are foraging. Ensure bees are not foraging in the crop to be treated. Remove beehives from areas due to be sprayed. Ensure alternative (untreated) source of nectar is available for bees.
AFFIRM INSECTICIDE	emamectin as benzoate(17g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AGCHEM CHLORPYRIFOS 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AGRIMEC MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AGSPRAY ABAMECTIN MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AGSPRAY APHID GUARD 200 SC APHICIDE/INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AGSPRAY CHLORPYRIFOS 500EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AKOSTAR INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
ALL-MITEY 290 SL INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower when bees are foraging.
ALPHA DUOP 100 INSECTICIDE	alpha-cypermethrin(100g/L)	DO NOT spray on any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
ALPHASIP DUO100 INSECTICIDE	alpha-cypermethrin(100g/L)	DO NOT spray on any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
AMBUSH EC INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AMBUSH EMULSIFIABLE CONCENTRATE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AMGROW FRUIT FLY CONTROL	spinosad(0.24g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
AMGROW SYSTEMIC INSECT & MITE SPRAY	dimethoate(0.3g/kg)	DO NOT spray if bees are feeding on flowering plants. Will kill bees.
APPARENT CHLORPYRIFOS 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ARRIVO 250 EC INSECTICIDE	cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
ARRIVO 40 ULV INSECTICIDE	cypermethrin(40g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
AVID MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AW AGFEN 100EC INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
AW ALF 100 EC INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
AW ANNOUNCE MITICIDE / INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AW CHOP 500 INSECTICIDE AND TERMITICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AW DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
AW FLAK INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.
AW MITEFEN 80 INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
AXE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BANSHEE EC INSECTICIDE	beta-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BARMAC DIAZINON INSECTICIDE	diazinon(800g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BARMAC OUT OF BOUNDS INSECTICIDE AND TERMITICIDE	bifenthrin(100g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
BAYTEX 550 INSECTICIDE SPRAY	fenthion(550g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BAYTHROID 50 EC INSECTICIDE	cyfluthrin(50g/L)	Dangerous to bees and other beneficial insects. DO NOT spray any plants in flower while bees are foraging.
BIOMECTIN MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BUGMASTER FLOWABLE INSECTICIDE	carbaryl(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BULLDOCK 25 EC INSECTICIDE	beta-cyfluthrin(25g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
BULLDOCK DUO INSECTICIDE	beta-cyfluthrin(25g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CAMPBELL ANTARES 100 EC INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CAMPBELL BENTHION 200 FLOWABLE INSECTICIDE	azinphos-methyl(200g/L)	Dangerous to bees. Will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift. Residues toxic to bees may remain for several days after application.
CAMPBELL BIFF 100SC INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CAMPBELL KAISER 200SC INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
CAMPBELL PENNSIDE FLOWABLE MICROENCAPSULATED INSECTICIDE	diazinon(240g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHEMAG BIFEN 100 INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CHEMAG BIFEN 80SC INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CHEMAG DIMETHOATE INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHEMAG METHOMYL 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
CHEMAG OMEN 290 INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHEMAG SENTINEL 800 INSECTICIDE SPRAY	omethoate(800g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHEMFORCE ALPHA-CYPERMETHRIN 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
CHEMFORCE CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHEMICIDE 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHIX EC INSECTICIDE	beta-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CHOICE BIFENDOFF 100 INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
COMPEL INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CONFIDOR 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CONFIDOR GARDEN INSECTICIDE	imidacloprid(50g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CONQUEST ALPHA DUO 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
CONQUEST ALPHA FORTE 250 SC INSECTICIDE	alpha-cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
CONQUEST ARROW 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CONQUEST CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CONQUEST CYPERMETHRIN 250 INSECTICIDE	cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
CONQUEST DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
COUNTRY ABAMECTIN INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
COUNTRY ALPHA-CYPERMETHRIN 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
COUNTRY BIFENTHRIN 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
COUNTRY BIFENTHRIN DUO 100 EC TERMITICIDE & INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
COUNTRY CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
COUNTRY DIAZINON 800 INSECTICIDE	diazinon(800g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
COURAZE 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CROP CARE DOMINEX DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
CROP CARE SECURE 360 SC INSECTICIDE-MITICIDE	chlorfenapyr(360g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CROPRO BUZZARD INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
CROPRO D-SECT EC INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
CROPRO POUNCE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.





# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
CROPRO STALK INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CROPRO STEALTH MITICIDE AND INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CROPRO ZEUS TERMITICIDE AND INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
CYAN PLANT GROWTH REGULATOR	cyanamide(530g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CYPERSHIELD ULV 40 INSECTICIDE	cypermethrin(40g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
CYREN 300 ULV/EC INSECTICIDE	chlorpyrifos(300g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CYREN 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CYREN 500 WP INSECTICIDE	chlorpyrifos(500g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CYREN PC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
CYRUX 200 EC INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
CYRUX 250 EC INSECTICIDE	cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
CYRUX 40 ULV INSECTICIDE	cypermethrin(40g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DANADIM INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Apply in the afternoon when bees have finished foraging.
DAVID GRAYS CARBARYL 500 FLOWABLE INSECTICIDE	carbaryl(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DAVID GRAYS CHLORPYRIFOS 500	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DAVID GRAYS CRICKET & GRASSHOPPER KILLER BAIT	carbaryl(18g/kg)	Dangerous to bees
DAVID GRAYS FENITROTHION 1000	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DAVID GRAYS LAWN BEETLE GRANULES	chlorpyrifos(40g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DAVID GRAYS PCO CHLORPYRIFOS 500	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DAVID GRAYS THERMAL FOGGING AND ULV INSECTICIDE CONCENTRATE	bioresmethrin(50g/L)+pip .but.(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DECIS 110 EC INSECTICIDE	deltamethrin(110g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
DECIS OPTIONS INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
DELEGATE INSECTICIDE	spinetoram(250g/kg)	Bee safety: Delegate is dangerous to bees and will kill bees foraging in the crop being treated or in hives which are oversprayed or reached by spray-drift. Once the spray deposit has dried, foraging bees should not be affected when using spray volumes of 2000 L/ha or less. However, if using spray volumes greater than 2000 L/ha, it is possible that foraging bees may be affected for some days after spraying.
DELTAGUARD ULV INSECTICIDE	deltamethrin(5.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
DELTASHIELD 27.5 INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
DELTASHIELD 5.5 INSECTICIDE	deltamethrin(5.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
DICTATE 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DICTATE ULV INSECTICIDE	alpha-cypermethrin(16g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DIPTEREX 500 SL INSECTICIDE	trichlorfon(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DISECT 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
DOMINEX 16 ULV INSECTICIDE	alpha-cypermethrin(16g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DOMINEX DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
DORMEX PLANT GROWTH REGULATOR	cyanamide(520g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
DUOMAX HC520 PLANT GROWTH REGULATOR	cyanamide(520g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
DUPONT AVATAR INSECTICIDE	indoxacarb(300g/kg)	Dangerous to bees. DO NOT apply when bees are actively foraging. Avoid direct application or drift of the spray mix onto beehives. After the spray has dried, bees can safely forage flowering crops.
DUPONT LANNATE L INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
DUPONT MARLIN INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
DUPONT STEWARD EC INSECTICIDE	indoxacarb(150g/L)	Dangerous to bees. DO NOT apply when bees are actively foraging. Avoid direct application or drift of the spray mix onto beehives. After the spray has dried, bees can safely forage flowering crops.
DURAMITE MITICIDE	bifenazate(480g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ECHEM ABAMECTIN INSECTICIDE/MITICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ECHEM ALPHA-CYP 100 DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
ECHEM ALPHA-CYP ULV INSECTICIDE	alpha-cypermethrin(16g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
E-CHEM CYPERMETHRIN 250 EC INSECTICIDE	cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: If flowering has started, application should be deferred until flowering is complete but before heads turn down. If treatment is unavoidable during flowering and bees are actively foraging in the crop, spraying must take place in the very early morning or very late afternoon.
ECHEM DELTAMETHRIN DUO INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
ECHEM DELTAMETHRIN ULV INSECTICIDE	deltamethrin(5.5g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
ENTRUST NATURALYTE INSECT CONTROL	spinosad(800g/kg)	Dangerous to bees. Avoid direct application or drift of the spray mix onto beehives. DO NOT spray on plants in flower if bees are active. Once the spray deposit has dried, foraging bees will not be affected.
ENVIROGUARD TURF & ORNAMENTAL RESIDUAL INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
ENVIROMAX BIFENTHRIN 80SC PROFESSIONAL INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
ENVIROMAX BIFENTHRIN 80SC RESIDUAL INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
ERASER 750 INSECTICIDE	acephate(750g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMALINX ABACIN 18 EC INSECTICIDE/MITICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMALINX ALPHACYPER INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
FARMALINX CHLORPOS INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMALINX IMI 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMALINX OMETHO-MITE 290 SL INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
FARMOZ ALPHA-SCUD ELITE INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
FARMOZ ALPHA-SCUD ULV INSECTICIDE	alpha-cypermethrin(16g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ BALLISTIC ELITE INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
FARMOZ CYREN 500 WP INSECTICIDE	chlorpyrifos(500g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ DIAZOL 800 INSECTICIDE	diazinon(800g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are actively foraging. DO NOT spray when bees are foraging. Spray in late afternoon after bees have finished foraging.
FARMOZ DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
FARMOZ ELECTRA 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
FARMOZ FENITROTHION 1000 INSECTICIDE	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ GUSATHION 200 SC INSECTICIDE	azinphos-methyl(200g/L)	Dangerous to bees. Will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift. Residues toxic to bees may remain for several days after application.
FARMOZ HELLFIRE 500 EC INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ INSTINCT 300 COTTON INSECTICIDE	chlorpyrifos(300g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ KOHINOR 200 INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
FARMOZ SCUD ELITE INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have a deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in early morning and late evening while bees are not foraging.
FARMOZ SHOWDOWN 375 INSECTICIDE	thiodicarb(375g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ SORCERER 18 MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ STRIKE-OUT 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ STRIKE-OUT 500 WP INSECTICIDE	chlorpyrifos(500g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ STRIKE-OUT PC TERMITICIDE AND INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ SUPRATHION 400 EC INSECTICIDE	methidathion(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FARMOZ VENOM 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
FARMOZ VENOM 80 SC INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower when bees are foraging. Spray in the night or early morning when bees are not actively foraging.
FARMOZ WIZARD 18 MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FASTAC DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
FASTAC ULV INSECTICIDE	alpha-cypermethrin(16g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
FASTAC XCEL INSECTICIDE	alpha-cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
FENSTAR 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
FENTHRIN INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
FICAM D INSECTICIDE DUST	bendiocarb(10g/kg)	Dangerous to bees. DO NOT allow this product to contact any plants in flower while bees are foraging.
FICAM W INSECTICIDE	bendiocarb(800g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FLORAMITE MITICIDE	bifenazate(480g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FOLIMAT 800 INSECTICIDE SPRAY	omethoate(800g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FORTUNE 500 MULTI-PURPOSE INSECTICIDE AND TERMITICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FRONTIER-P HERBICIDE	dimethenamid-P(720g/L)	Toxic to bees.
FURY 14 ULV INSECTICIDE	zeta-cypermethrin(14g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FYFANON 440 EW INSECTICIDE	maldison(440g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FYFANON 500 EC INSECTICIDE	maldison(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
FYFANON ULV INSECTICIDE	maldison(1169g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT spray over or upwind of beehives. Remove bees from area before blanket spraying.
GENERIFOS 500EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.





# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
GENFARM ABA 18 MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.
GENFARM ALPHA DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
GENFARM BIFENTHRIN 100 MITICIDE/INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower when bees are foraging. Spray in the early morning when bees are not actively foraging.
GENFARM BOOM 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
GENFARM CENTAUR 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
GENFARM CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
GENFARM CYPER PLUS 250 INSECTICIDE	cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
GOOGLY ALPHA-DUO 250SC INSECTICIDE	alpha-cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
GRANULAR PRODUCTS IMIDACLOPRID 350 SOIL INSECTICIDE	imidacloprid(350g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
GRASS VALLEY ALPHA- CYPERMETHRIN 100 INSECTICIDE	alpha- cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
GREEN GUARD SC BIOLOGICAL INSECTICIDE	M. anisopliae var. acridum(40000Msp/g)	This product is a potential hazard to bees. DO NOT spray on plants in flower while bees are foraging.
GREEN GUARD SC PREMIUM BIOLOGICAL INSECTICIDE	M. anisopliae var. acridum(100g/L)	This product is a potential hazard to bees. DO NOT spray on plants in flower while bees are foraging.
GREMLIN MITICIDE AND INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
GRINGO INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
HALLEY ALPHA- CYPERMETHRIN 100 INSECTICIDE	alpha- cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
HALLEY CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
HALLEY CYPERMETHRIN 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. This product is highly toxic to bees and should not be applied while bees are actively foraging.
HALLEY DELTAMETHRIN DUO INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
HALLEY DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Apply in the afternoon when bees have finished foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
HY-MAL INSECTICIDE	maldison(1150g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT spray over or upwind of beehives. Remove bees from area before blanket spraying.
IMIDAN INSECTICIDE/MITICIDE	phosmet(150g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
IMTRADE ABACHEM 18 MITICIDE/INSECTICIDE	abamectin(18g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
IMTRADE ANNIHILATE 200 SC ALL PURPOSE INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
IMTRADE CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
IMTRADE CONFRONT 800 WG INSECTICIDE	thiodicarb(800g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
IMTRADE CYPERSHIELD 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
IMTRADE DELTA-DUO INSECTICIDE	deltamethrin(27.5g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
IMTRADE DICTATE DUO 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
IMTRADE KUNG FU 250 INSECTICIDE	lambda-cyhalothrin(250g/L)	Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.
IMTRADE OMEN 290 INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
INNOVA ALPHA 100 DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
INNOVA BIFENTHRIN 100 MITICIDE / INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
INTREPID 360 SC INSECTICIDE-MITICIDE	chlorfenapyr(360g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KARATE WITH ZEON TECHNOLOGY INSECTICIDE	lambda-cyhalothrin(250g/L)	Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.
KDPC PY-OMNI INSECTICIDE WITH NATURAL PYRETHRUM	pyrethrins(40g/L)+pip.but .(160g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KEN-CYPER 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
KENDON CARBARYL WETTABLE POWDER INSECTICIDE	carbaryl(800g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KENDON FENITROTHION 1000EC INSECTICIDE	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KENDON PERMEKIL EMULSIFIABLE CONCENTRATE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KENSO AGCARE KENSBAN 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
KENSO AGCARE KEN-TAC 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
KENSO AGCARE SAVAGE 200 INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
KENSO AGCARE TAL-KEN 100 INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
KILL-A-MITE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LANCER 750 DF INSECTICIDE	acephate(750g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LANCER 750 SP INSECTICIDE	acephate(750g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LANCER 970 INSECTICIDE	acephate(970g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LANNATE-L INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
LARVIN 375 INSECTICIDE	thiodicarb(375g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LARVIN 800 WG INSECTICIDE	thiodicarb(800g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LEBAYCID FRUIT FLY & INSECT KILLER	fenthion(80g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LEBAYCID INSECTICIDE SPRAY	fenthion(550g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
LEGION 200SC INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.
LE-MAT 290 SL INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LORSBAN 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
LORSBAN 750 WG INSECTICIDE	chlorpyrifos(750g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MACPHERSONS SENECA INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
MASPART IMI-FLOW 200 INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MASTERGUARD INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
MATADOR WITH ZEON TECHNOLOGY INSECTICIDE	lambda-cyhalothrin(250g/L)	Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.
MERIT TURF AND ORNAMENTAL INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MESUROL 500 SC INSECTICIDE	methiocarb(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
MESUROL 750 BIRD REPELLENT AND SNAIL AND SLUG SPRAY	methiocarb(750g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
METHYLATE 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
MILBEKNOCK MITICIDE	milbemectin(9.3g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MITE MASTER 290 INSECTICIDE	omethoate(290g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MITE TERMINATOR	abamectin(18g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
MONITOR INSECTICIDE	methamidophos(580g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
MUSTANG INSECTICIDE	ethion(360g/L)+zeta-cypermethrin(20g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
NATURALURE FRUIT FLY BAIT CONCENTRATE	spinosad(0.24g/L)	Dangerous to bees. Avoid direct application or drift of the spray mix onto beehives. DO NOT spray on plants in flower if bees are active. Once the spray deposit has dried, foraging bees will not be affected.
NEVWEB FENITROTHION 1000 INSECTICIDE	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NITOFOL INSECTICIDE SPRAY	methamidophos(580g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUDRIN 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
NUFARM ABAMECT INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
NUFARM ADONIS 3UL INSECTICIDE	fipronil(3g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.
NUFARM ASTOUND DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
NUFARM CHLORPYRIFOS 500 EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM CHLORPYRIFOS PCO INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM DIMETHOATE SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM FASTAC DUO INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
NUFARM FENITROTHION 1000 INSECTICIDE	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM LEPIDEX 500 INSECTICIDE	trichlorfon(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.





# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
NUFARM MALDISON 500 INSECTICIDE	maldison(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM NUPRID 200SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
NUFARM NUPRID 350SC INSECTICIDE	imidacloprid(350g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.
NUFARM REGENT 200SC INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.
ORTHENE XTRA INSECTICIDE	acephate(970g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
OSPRAY ALPHA-CYPERMETHRIN 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
OSPRAY COURAZE 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
OSPRAY COURAZE CLASSIC INSECTICIDE	imidacloprid(350g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
OSPRAY METHOMYL 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
OSPRAY PERMETHRIN 500 INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
OSPRAY TROJAN INSECTICIDE	gamma-cyhalothrin(150g/L)	Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.
PACIFIC IMIDACLOPRID 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PEGASUS MITICIDE/INSECTICIDE	diafenthiuron(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PERMERID 500 EC INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PHOSDRIN INSECTICIDE	mevinphos(1110g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PIDGEON'S PEST CONTROLLER 500 TERMITICIDE AND INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
POUNCE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PROCLAIM INSECTICIDE	emamectin as benzoate(44g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PROVADO 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PROWLER 300 INSECTICIDE	chlorpyrifos(300g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PY-BO NATURAL PYRETHRUM INSECTICIDAL CONCENTRATE	pyrethrins(80g/L)+pip.but .(480g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PYGANIC ORGANIC INSECTICIDE	pyrethrins(13g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
PYRINEX 250 WP INSECTICIDE	chlorpyrifos(250g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
PY-ZAP INSECTICIDE WITH NATURAL PYRETHRUM	pyrethrins(40g/L)+pip.but .(160g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
RAINBOW ABAMECTIN INSECTICIDE / MITICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
RAINBOW CHLORPYRIFOS 500 INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
REGENT 200SC INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
REGENT 800WG INSECTICIDE	fipronil(800g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ROGOR UPGRADE INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ROTAM ACARMIK INSECTICIDE/MITICIDE EC	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
ROVER SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
RYGEL ALPHA FORTE 250 SC INSECTICIDE	alpha-cypermethrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
RYGEL IMIDACLOPRID 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SABERO CHLORPYRIFOS 500EC INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SABOTEUR SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SANMITE MITICIDE	pyridaben(250g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SARRITOR INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower when bees are foraging. Spray in the early morning when bees are not actively foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
SCOTTS MAXGUARD 80SC INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
SCOTTS PROCIDE 80SC INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
SECURE 360 SC INSECTICIDE-MITICIDE	chlorfenapyr(360g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SENATOR 200 SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SENTINEL 80SC INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
SILVASHIELD INJECTABLE TREE INSECTICIDE	imidacloprid(200g/L)	This product is highly toxic to bees exposed to direct treatment or residues on flowering trees and shrubs. DO NOT use on trees which are likely to be used by commercial beehives.
SINMAS 225 INSECTICIDE	methomyl(225g/L)	Dangerous to bees. DO NOT spray any plants or flowers while bees are foraging. Ensure beehives are removed from the area to be treated and from adjacent paddocks.
SIPCAM PYRANICA MITICIDE	tebufenpyrad(200g/kg)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SONIC 200EC INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
STAKEOUT EMULSIFIABLE CONCENTRATE INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUCCESS NEO INSECTICIDE	spinetoram(120g/L)	This product is highly toxic to bees; will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift. Residues may remain toxic to bees for several days after application.
SUCCESS2 NATURALYTE INSECT CONTROL	spinosad(240g/L)	Dangerous to bees. Avoid direct application or drift of the spray mix onto beehives.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
SUMI-ALPHA FLEX INSECTICIDE	esfenvalerate(50g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUMITHION 1000EC INSECTICIDE	fenitrothion(1000g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUMITOMO SAMURAI SYSTEMIC INSECTICIDE	clothianidin(500g/kg)	Dangerous to bees and will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift, and residues may remain toxic to bees several days after application. Toxic to bees for several days after application.
SUMITOMO SHIELD SYSTEMIC INSECTICIDE	clothianidin(200g/L)	Dangerous to bees and will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift, and residues may remain toxic to bees several days after application. Toxic to bees for several days after application. Used on eucalypt seedlings to 1 year old.
SUMITOMO STEALTH SYSTEMIC INSECTICIDE	clothianidin(500g/kg)	Dangerous to bees and will kill bees foraging in the crop to be treated or in hives which are oversprayed or reached by spray drift, and residues may remain toxic to bees several days after application. Toxic to bees for several days after application.
SUMITOMO SUMITHION ULV PREMIUM GRADE INSECTICIDE	fenitrothion(1.23kg/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUPERWAY DIMETHOATE 300 SYSTEMIC INSECTICIDE	dimethoate(300g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
SUPERWAY DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUPRACIDE 400 EMULSIFIABLE CONCENTRATE INSECTICIDE	methidathion(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUREFIRE FIVESTAR INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower when bees are foraging. Spray in the night or early morning when bees are not actively foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
SUREFIRE ORCHARD & GARDEN INSECTICIDE	dimethoate(300g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
SUREFIRE SPECTRUM 200SC INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
SUREFIRE VISTA 200SC INSECTICIDE	fipronil(200g/L)	Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or expected to flower within 28 days (7 days pasture and sorghum). Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds, or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the registration number, for assistance in contacting the owner.
TALSTAR 100 EC INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
TALSTAR 250 EC INSECTICIDE/MITICIDE	bifenthrin(250g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
TALSTAR 80 SC INSECTICIDE/MITICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
TITAN ALPHA DUO 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
TITAN BIFENTHRIN 100 INSECTICIDE/ MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
TITAN CHLORPYRIFOS 500 TERMITICIDE AND INSECTICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
TITAN CYPERMETHRIN 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Sunflowers: Note: once flowering has commenced, spraying should be delayed until after flowering has ceased but before heads turn down to avoid damage to foraging bees. If treatment is unavoidable due to pest numbers apply during very early morning or very late afternoon when bee activity is minimal.
TITAN DIMETHOATE 400 SYSTEMIC INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Some repellent effect may be apparent for approximately 2 days.
TITAN IMIDACLOPRID 200 INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
TROJAN INSECTICIDE	gamma-cyhalothrin(150g/L)	Toxic to bees. DO NOT spray when bees are actively foraging. Risk is reduced by spraying in the early morning or late evening.
TURF CULTURE CEASEFIRE 80 SC INSECTICIDE	bifenthrin(80g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the night or early morning when bees are not actively foraging.
UNITED FARMERS BIFENTHRIN 100 INSECTICIDE/MITICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
UNITED FARMERS CHLORPYRIFOS 500 INSECTICIDE AND TERMITICIDE	chlorpyrifos(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
UNITED FARMERS CYPERMETHRIN 200 INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.



# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
UNITED FARMERS UNIALPHACYPER 100 INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. To protect bees and ensure adequate pollination of sunflowers, application during flowering should be avoided. If application is necessary at flowering apply early morning or late afternoon when bees are not actively foraging.
UNITED FARMERS UNIDIME 400 INSECTICIDE	dimethoate(400g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
UNITOX 100 EC INSECTICIDE	alpha-cypermethrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. This product is known to have deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in the early morning and late evening when bees are not foraging.
VANTAL 18 EC MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
VANTAL 18 EW MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
VERTIMEC MITICIDE/INSECTICIDE	abamectin(18g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
VOODOO 200 INSECTICIDE	imidacloprid(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
WEBCOT S-PY NATURAL PYRETHRUM INSECTICIDE	pyrethrins(40g/L)+pip.but .(160g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.
WEBZONE TERMITICIDE & INSECTICIDE	bifenthrin(100g/L)	Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.
WSD CHLORPYRIFOS 500 EC INSECTICIDE	chlorpyrifos(500g/L)	No bee warning statement on the label. This product has been included on the basis that it has the same active constituent(s), active constituent(s) concentration, application rate and intended use as other products which do contain a bee related warning on the label.
WSD CYPERMETHRIN 200 EC INSECTICIDE	cypermethrin(200g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.





# Appendix 2



Product name	Active constituent(s) and concentration	Bee related label statement
ZEEPER 500 EC INSECTICIDE	permethrin 40:60(500g/L)	Dangerous to bees. DO NOT spray any plants in flower where bees are foraging.

